arXiv:1010.5688v1 [math.FA] 27 Oct 2010

THE TOPOLOGICAL CENTERS AND FACTORIZATION PROPERTIES OF MODULE ACTIONS AND $*-INVOLUTION$ ALGEBRAS

KAZEM HAGHNEJAD AZAR

Abstract. For Banach left and right module actions, we extend some propositions from Lau and *Ülger* into general situations and we establish the relationships between topological centers of module actions. We also introduce the new concepts as Lw^*w-property and Rw^*w-property for Banach $A-bimodule$ B and we obtain some conclusions in the topological center of module actions and Arens regularity of Banach algebras. we also study some factorization properties of left module actions and we find some relations of them and topological centers of module actions. For Banach algebra A, we extend the definition of $*-involution$ algebra into Banach $A-bimodule$ B with some results in the factorizations of B^*. We have some applications in group algebras.

1.Introduction and Preliminaries

As is well-known [1], the second dual A^{**} of A endowed with the either Arens multiplications is a Banach algebra. The constructions of the two Arens multiplications in A^{**} lead us to definition of topological centers for A^{**} with respect both Arens multiplications. The topological centers of Banach algebras, module actions and applications of them were introduced and discussed in [6, 8, 13, 14, 15, 16, 17, 21, 22], and they have attracted by some attentions.

Now we introduce some notations and definitions that we used throughout this paper. Let A be a Banach algebra. We say that a net $(e_\alpha)_{\alpha\in I}$ in A is a left approximate identity ($=LAI$) [resp. right approximate identity ($=RAI$)] if, for each $a\in A$, $e_\alpha a\longrightarrow a$ [resp. $ae_\alpha\longrightarrow a$]. For $a\in A$ and $a'\in A^*$, we denote by $a'a$ and aa' respectively, the functionals on A^* defined by $\langle a'a,b\rangle=\langle a',ab\rangle=a'(ab)$ and $\langle aa',b\rangle=\langle a',ba\rangle=a'(ba)$ for all $b\in A$. The Banach algebra A is embedded in its second dual via the identification $\langle a,a'\rangle$ - $\langle a',a\rangle$ for every $a\in A$ and $a'\in A^*$. We denote the set $\{a'a:\ a\in A\ and\ a'\in A^*\}$ and $\{aa':\ a\in A\ and\ a'\in A^*\}$ by A^*A and AA^*, respectively, clearly these two sets are subsets of A^*. Let A has a BAI. If the equality $A^*A=A^*$, ($AA^*=A^*$) holds, then we say that A^* factors on the left (right). If both equalities $A^*A=AA^*=A^*$ hold, then we say that A^* factors on both sides. Let X,Y,Z be normed spaces and $m:X\times Y\to Z$ be a bounded bilinear mapping. Arens in [1] offers two natural extensions m^{***} and m^{t***t} of m from $X^{**}\times Y^{**}$ into Z^{**} as following:

1. $m^*:Z^*\times X\to Y^*$, given by $\langle m^*(z',x),y\rangle=\langle z',m(x,y)\rangle$ where $x\in X$, $y\in Y$,

---

2000 *Mathematics Subject Classification.* 46L06; 46L07; 46L10; 47L25.

Key words and phrases. Arens regularity, bilinear mappings, Topological center, Second dual, Module action, factorization, $*-involution$ algebra.

$z' \in Z^*$,
2. $m^{**} : Y^{**} \times Z^* \to X^*$, given by $\langle m^{**}(y'', z'), x\rangle = \langle y'', m^*(z', x)\rangle$ where $x \in X$, $y'' \in Y^{**}$, $z' \in Z^*$,
3. $m^{***} : X^{**} \times Y^{**} \to Z^{**}$, given by $\langle m^{***}(x'', y''), z'\rangle = \langle x'', m^{**}(y'', z')\rangle$
where $x'' \in X^{**}$, $y'' \in Y^{**}$, $z' \in Z^*$.
The mapping m^{***} is the unique extension of m such that $x'' \to m^{***}(x'', y'')$ from X^{**} into Z^{**} is $weak^* - to - weak^*$ continuous for every $y'' \in Y^{**}$, but the mapping $y'' \to m^{***}(x'', y'')$ is not in general $weak^* - to - weak^*$ continuous from Y^{**} into Z^{**} unless $x'' \in X$. Hence the first topological center of m may be defined as following

$$Z_1(m) = \{x'' \in X^{**} : \; y'' \to m^{***}(x'', y'') \; is \; weak^* - to - weak^* - continuous\}.$$

Let now $m^t : Y \times X \to Z$ be the transpose of m defined by $m^t(y, x) = m(x, y)$ for every $x \in X$ and $y \in Y$. Then m^t is a continuous bilinear map from $Y \times X$ to Z, and so it may be extended as above to $m^{t***} : Y^{**} \times X^{**} \to Z^{**}$. The mapping $m^{t***t} : X^{**} \times Y^{**} \to Z^{**}$ in general is not equal to m^{***}, see [1], if $m^{***} = m^{t***t}$, then m is called Arens regular. The mapping $y'' \to m^{t***t}(x'', y'')$ is $weak^* - to - weak^*$ continuous for every $y'' \in Y^{**}$, but the mapping $x'' \to m^{t***t}(x'', y'')$ from X^{**} into Z^{**} is not in general $weak^* - to - weak^*$ continuous for every $y'' \in Y^{**}$. So we define the second topological center of m as

$$Z_2(m) = \{y'' \in Y^{**} : \; x'' \to m^{t***t}(x'', y'') \; is \; weak^* - to - weak^* - continuous\}.$$

It is clear that m is Arens regular if and only if $Z_1(m) = X^{**}$ or $Z_2(m) = Y^{**}$. Arens regularity of m is equivalent to the following

$$\lim_i \lim_j \langle z', m(x_i, y_j)\rangle = \lim_j \lim_i \langle z', m(x_i, y_j)\rangle,$$

whenever both limits exist for all bounded sequences $(x_i)_i \subseteq X$, $(y_i)_i \subseteq Y$ and $z' \in Z^*$, see [6, 18].
The regularity of a normed algebra A is defined to be the regularity of its algebra multiplication when considered as a bilinear mapping. Let a'' and b'' be elements of A^{**}, the second dual of A. By *Goldstin's* Theorem [6, P.424-425], there are nets $(a_\alpha)_\alpha$ and $(b_\beta)_\beta$ in A such that $a'' = weak^* - \lim_\alpha a_\alpha$ and $b'' = weak^* - \lim_\beta b_\beta$. So it is easy to see that for all $a' \in A^*$,

$$\lim_\alpha \lim_\beta \langle a', m(a_\alpha, b_\beta)\rangle = \langle a''b'', a'\rangle$$

and

$$\lim_\beta \lim_\alpha \langle a', m(a_\alpha, b_\beta)\rangle = \langle a''ob'', a'\rangle,$$

where $a''b''$ and $a''ob''$ are the first and second Arens products of A^{**}, respectively, see [6, 14, 18].
The mapping m is left strongly Arens irregular if $Z_1(m) = X$ and m is right strongly Arens irregular if $Z_2(m) = Y$.
This paper is organized as follows.
a) In section two, for a Banach $A - bimodule$, we have

(1) $a'' \in Z_{B^{**}}(A^{**})$ if and only if $\pi_\ell^{****}(b', a'') \in B^*$ for all $b' \in B^*$.
(2) $F \in Z_{B^{**}}((A^*A)^*)$ if and only if $\pi_\ell^{****}(g, F) \in B^*$ for all $g \in B^*$.
(3) $G \in Z_{(A^*A)^*}(B^{**})$ if and only if $\pi_r^{****}(g, G) \in A^*A$ for all $g \in B^*$.

(4) Let B has a BAI $(e_\alpha)_\alpha \subseteq A$ such that $e_\alpha \overset{w^*}{\to} e''$. Then if $Z^t_{e^{**}}(B^{**}) = B^{**}$ [resp. $Z_{e^{**}}(B^{**}) = B^{**}$] and B^* factors on the left [resp. right], but not on the right [resp. left], then $Z_{B^{**}}(A^{**}) \neq Z^t_{B^{**}}(A^{**})$.

(5) $B^*A \subseteq wap_\ell(B)$ if and only if $AA^{**} \subseteq Z_{B^{**}}(A^{**})$.

(6) Let $b' \in B^*$. Then $b' \in wap_\ell(B)$ if and only if the adjoint of the mapping $\pi^*_\ell(b', \;) : A \to B^*$ is $weak^* - to - weak$ continuous.

Then $(\alpha) \Rightarrow (\beta) \Rightarrow (\gamma)$.

If we take $T \in LM(A, B)$ and if B has a sequential BAI and it is WSC, then (α), (β) and (γ) are equivalent.

b) In section three, for a Banach $A-bimodule\ B$, we define $Left-weak^*-to-weak$ property [$=Rw^*w-$ property] and $Right - weak^* - to - weak$ property [$=Rw^*w-$ property] for Banach algebra A and we show that

(1) If $A^{**} = a_0A^{**}$ [resp. $A^{**} = A^{**}a_0$] for some $a_0 \in A$ and a_0 has Rw^*w- property [resp. Lw^*w- property], then $Z_{B^{**}}(A^{**}) = A^{**}$.

(2) If $B^{**} = a_0B^{**}$ [resp. $B^{**} = B^{**}a_0$] for some $a_0 \in A$ and a_0 has Rw^*w- property [resp. Lw^*w- property] with respect to B, then $Z_{A^{**}}(B^{**}) = B^{**}$.

(3) If B^* factors on the left [resp. right] with respect to A and A has Rw^*w- property [resp. Lw^*w- property], then $Z_{B^{**}}(A^{**}) = A^{**}$.

(4) If B^* factors on the left [resp. right] with respect to A and A has Rw^*w- property [resp. Lw^*w- property] with respect B, then $Z_{A^{**}}(B^{**}) = B^{**}$.

(5) If $a_0 \in A$ has Rw^*w- property with respect to B, then $a_0A^{**} \subseteq Z_{B^{**}}(A^{**})$ and $a_0B^* \subseteq wap_\ell(B)$.

(6) Assume that $AB^* \subseteq wap_\ell B$. If B^* strong factors on the left [resp. right], then A has Lw^*w- property [resp. Rw^*w- property] with respect to B.

(7) Assume that $AB^* \subseteq wap_\ell B$. If B^* strong factors on the left [resp. right], then A has Lw^*w- property [resp. Rw^*w- property] with respect to B.

c) In section four, for a left Banach $A - module\ B$, we study some relationships between factorization properties of left module action and topological centers of it.

(1) Let $(e_\alpha)_\alpha \subseteq A$ be a $LBAI$ for B. Then the following assertions hold.

i) For each $b' \in B^*$, $\pi^*_\ell(b', e_\alpha) \overset{w^*}{\to} b'$.

ii) B^* factors on the left with respect to A if and only if B^{**} has a W^*LBAI $(e_\alpha)_\alpha \subseteq A$.

iii) B^{**} has a W^*LBAI $(e_\alpha)_\alpha \subseteq A$ if and only if B^{**} has a left unit element $e'' \in A^{**}$ such that $e_\alpha \overset{w^*}{\to} e''$.

(2) Suppose that $b' \in wap_\ell(B)$. Let $a'' \in A^{**}$ and $(a_\alpha)_\alpha \subseteq A$ such that $a_\alpha \overset{w^*}{\to} a''$ in A^{**}. Then we have

$$\pi^*_\ell(b', a_\alpha) \overset{w}{\to} \pi^{****}_\ell(b', a'').$$

(3) Let B^* factors on the left with respect to A. If $AA^{**} \subseteq Z_{B^{**}}(A^{**})$, then $Z_{B^{**}}(A^{**}) = A^{**}$.

(4) Let B^{**} has a $LBAI$ with respect to A^{**}. Then B^{**} has a left unit with respect to A^{**}.

(5) Let B has a $LBAI$ with respect to A. Then we have the following assertions.

i) B^* factors on the left with respect to A if and only if for each $b' \in B^*$, we have $\pi_\ell^*(b', e_\alpha) \overset{w}{\to} b'$ in B^*.

ii) B factors on the left with respect to A if and only if for each $b \in B$, we have $\pi_\ell^*(b, e_\alpha) \overset{w}{\to} b$ in B.

(6) Let A has a $LBAI$ $(e_\alpha)_\alpha \subseteq A$ such that $e_\alpha \overset{w^*}{\to} e''$ in A^{**} where e'' is a left unit for A^{**}. Suppose that $Z^t_{e''}(B^{**}) = B^{**}$. Then, B factors on the right with respect to A if and only if e'' is a left unit for B^{**}.

d) In section five, for a Banach $A-bimodule$ B we study $*-involution$ algebra on B^{**} with respect to first Arens product with some results in the factorization in B^*, that is, suppose that $(e_\alpha)_\alpha \subseteq A$ is a BAI for B. Then if B^{**} is a Banach $*-involution$ algebra as A^{**}-module, then B^{**} is unital as A^{**}-module and B^* factors on the both side.

2. The topological centers of module actions

Let B be a Banach $A-bimodule$, and let

$$\pi_\ell : \ A \times B \to B \ and \ \pi_r : \ B \times A \to B.$$

be the left and right module actions of A on B. Then B^{**} is a Banach $A^{**}-bimodule$ with module actions

$$\pi_\ell^{***} : \ A^{**} \times B^{**} \to B^{**} \ and \ \pi_r^{***} : \ B^{**} \times A^{**} \to B^{**}.$$

Similarly, B^{**} is a Banach $A^{**}-bimodule$ with module actions

$$\pi_\ell^{t***t} : \ A^{**} \times B^{**} \to B^{**} \ and \ \pi_r^{t***t} : \ B^{**} \times A^{**} \to B^{**}.$$

We may therefore define the topological centers of the right and left module actions of A on B as follows:

$$Z_{A^{**}}(B^{**}) = Z(\pi_r) = \{b'' \in B^{**} : \ the \ map \ a'' \to \pi_r^{***}(b'', a'') \ : \ A^{**} \to B^{**}$$
$$is \ weak^* - to - weak^* \ continuous\}$$
$$Z_{B^{**}}(A^{**}) = Z(\pi_\ell) = \{a'' \in A^{**} : \ the \ map \ b'' \to \pi_\ell^{***}(a'', b'') \ : \ B^{**} \to B^{**}$$
$$is \ weak^* - to - weak^* \ continuous\}$$
$$Z^t_{A^{**}}(B^{**}) = Z(\pi_\ell^t) = \{b'' \in B^{**} : \ the \ map \ a'' \to \pi_\ell^{t***}(b'', a'') \ : \ A^{**} \to B^{**}$$
$$is \ weak^* - to - weak^* \ continuous\}$$
$$Z^t_{B^{**}}(A^{**}) = Z(\pi_r^t) = \{a'' \in A^{**} : \ the \ map \ b'' \to \pi_r^{t***}(a'', b'') \ : \ B^{**} \to B^{**}$$
$$is \ weak^* - to - weak^* \ continuous\}$$

We note also that if B is a left(resp. right) Banach $A-module$ and $\pi_\ell : \ A \times B \to B$ (resp. $\pi_r : \ B \times A \to B$) is left (resp. right) module action of A on B, then B^* is a right (resp. left) Banach $A-module$.

We write $ab = \pi_\ell(a, b)$, $ba = \pi_r(b, a)$, $\pi_\ell(a_1a_2, b) = \pi_\ell(a_1, a_2b)$, $\pi_r(b, a_1a_2) = \pi_r(ba_1, a_2)$, $\pi_\ell^*(a_1b', a_2) = \pi_\ell^*(b', a_2a_1)$, $\pi_r^*(b'a, b) = \pi_r^*(b', ab)$, for all $a_1, a_2, a \in A$, $b \in B$ and $b' \in B^*$ when there is no confusion.

Theorem 2-1. We have the following assertions.

1. Assume that B is a Banach left $A-module$. Then, $a'' \in Z_{B^{**}}(A^{**})$ if and only if $\pi_\ell^{****}(b', a'') \in B^*$ for all $b' \in B^*$.
2. Assume that B is a Banach right $A-module$. Then, $b'' \in Z_{A^{**}}(B^{**})$ if and only if $\pi_r^{****}(b', b'') \in A^*$ for all $b' \in B^*$.

Proof. (1) Let $b'' \in B^{**}$. Then, for every $a'' \in Z_{B^{**}}(A^{**})$, we have

$$\langle \pi_\ell^{****}(b', a''), b'' \rangle = \langle b', \pi_\ell^{***}(a'', b'') \rangle = \langle \pi_\ell^{***}(a'', b''), b' \rangle$$

$$= \langle \pi_\ell^{t***t}(a'', b''), b' \rangle = \langle \pi_\ell^{t***}(b'', a''), b' \rangle = \langle b'', \pi_\ell^{t**}(a'', b') \rangle.$$

It follow that $\pi_\ell^{****}(b', a'') = \pi_\ell^{t**}(a'', b') \in B^*$.
Conversely, let $a'' \in A^{**}$ and let $\pi_\ell^{****}(a'', b') \in B^*$ for all $b' \in B^*$. Let $(b''_\alpha)_\alpha \subseteq B^{**}$ such that $b''_\alpha \overset{w^*}{\to} b''$. Then for every $b' \in B^*$, we have

$$\langle \pi_\ell^{***}(a'', b''_\alpha), b' \rangle = \langle b', \pi_\ell^{***}(a'', b''_\alpha) \rangle = \langle \pi_\ell^{****}(b', a''), b''_\alpha \rangle = \langle b''_\alpha, \pi_\ell^{****}(b', a'') \rangle$$

$$\to \langle b'', \pi_\ell^{****}(b', a'') \rangle = \langle \pi_\ell^{****}(b', a''), b'' \rangle = \langle \pi_\ell^{***}(a'', b''), b' \rangle.$$

Consequently $a'' \in Z_{B^{**}}(A^{**})$.

(2) Proof is similar to (1).

□

Theorem 2-2. Assume that B is a Banach $A-bimodule$. Then we have the following assertions.

1. $F \in Z_{B^{**}}((A^*A)^*)$ if and only if $\pi_\ell^{****}(g, F) \in B^*$ for all $g \in B^*$.
2. $G \in Z_{(A^*A)^*}(B^{**})$ if and only if $\pi_r^{****}(g, G) \in A^*A$ for all $g \in B^*$.

Proof. (1) Let $F \in Z_{B^{**}}((A^*A)^*)$ and $(b''_\alpha)_\alpha \subseteq B^{**}$ such that $b''_\alpha \overset{w^*}{\to} b''$. Then for all $g \in B^*$, we have

$$\langle \pi_\ell^{****}(g, F), b''_\alpha \rangle = \langle g, \pi_\ell^{***}(F, b''_\alpha) \rangle = \langle \pi_\ell^{***}(F, b''_\alpha), g \rangle$$

$$\to \langle \pi_\ell^{***}(F, b''), g \rangle = \langle \pi_\ell^{****}(g, F), b'' \rangle.$$

Thus, we conclude that $\pi_\ell^{****}(g, F) \in (B^{**}, weak^*)^* = B^*$.
Conversely, let $\pi_\ell^{****}(g, F) \in B^*$ for $F \in (A^*A)^*$ and $g \in B^*$. Assume that $b'' \in B^{**}$ and $(b''_\alpha)_\alpha \subseteq B^{**}$ such that $b''_\alpha \overset{w^*}{\to} b''$. Then

$$\langle \pi_\ell^{***}(F, b''_\alpha), g \rangle = \langle g, \pi_\ell^{***}(F, b''_\alpha) \rangle = \langle \pi_\ell^{****}(g, F), b''_\alpha \rangle$$

$$= \langle b''_\alpha, \pi_\ell^{****}(g, F) \rangle \to \langle b'', \pi_\ell^{****}(g, F) \rangle = \langle \pi_\ell^{****}(g, F), b'' \rangle$$

$$= \langle \pi_\ell^{***}(F, b''), g \rangle.$$

It follow that $F \in Z_{B^{**}}((A^*A)^*)$.

(2) Proof is similar to (1).

□

In the preceding theorems, if we take $B = A$, we obtain some parts of Lemma 3.1 from [14].

An element e'' of A^{**} is said to be a mixed unit if e'' is a right unit for the first Arens multiplication and a left unit for the second Arens multiplication. That is, e'' is a mixed unit if and only if for each $a'' \in A^{**}$, $a''e'' = e''oa'' = a''$. By [4, p.146], an element e'' of A^{**} is mixed unit if and only if it is a *weak** cluster point of some BAI $(e_\alpha)_{\alpha\in I}$ in A.
Let B be a Banach $A-$*bimodule* and $a'' \in A^{**}$. We define the locally topological center of the left and right module actions of a'' on B, respectively, as follows

$$Z^t_{a''}(B^{**}) = Z^t_{a''}(\pi^t_\ell) = \{b'' \in B^{**} :\ \pi_\ell^{t***t}(a'', b'') = \pi_\ell^{***}(a'', b'')\},$$
$$Z_{a''}(B^{**}) = Z_{a''}(\pi^t_r) = \{b'' \in B^{**} :\ \pi_r^{t***t}(b'', a'') = \pi_r^{***}(b'', a'')\}.$$

Thus we have

$$\bigcap_{a''\in A^{**}} Z^t_{a''}(B^{**}) = Z^t_{A^{**}}(B^{**}) = Z(\pi^t_r),$$
$$\bigcap_{a''\in A^{**}} Z_{a''}(B^{**}) = Z_{A^{**}}(B^{**}) = Z(\pi_r).$$

Let B be a Banach $A-$*bimodule*. We say that B is a left [resp. right] factors with respect to A, if $BA = B$ [resp. $AB = B$].

Definition 2-3. Let B be a Banach left $A-$*module* and $e'' \in A^{**}$ be a mixed unit for A^{**}. We say that e'' is a left mixed unit for B^{**}, if

$$\pi_\ell^{***}(e'', b'') = \pi_\ell^{t***t}(e'', b'') = b'',$$

for all $b'' \in B^{**}$.
The definition of right mixed unit for B^{**} is similar. B^{**} has a mixed unit, if it has equal left and right mixed unit.
It is clear that if $e'' \in A^{**}$ is a left (resp. right) unit for B^{**} and $Z_{e''}(B^{**}) = B^{**}$, then e'' is left (resp. right) mixed unit for B^{**}.

Theorem 2-4. Let B be a Banach $A-$*bimodule* with a *BAI* $(e_\alpha)_\alpha$ such that $e_\alpha \overset{w^*}{\to} e''$. Then if $Z^t_{e''}(B^{**}) = B^{**}$ [resp. $Z_{e''}(B^{**}) = B^{**}$] and B^* factors on the left [resp. right], but not on the right [resp. left], then $Z_{B^{**}}(A^{**}) \neq Z^t_{B^{**}}(A^{**})$.

Proof. Suppose that B^* factors on the left with respect to A, but not on the right. Let $(e_\alpha)_\alpha \subseteq A$ be a BAI for A such that $e_\alpha \overset{w^*}{\to} e''$. Thus for all $b' \in B^*$ there are $a \in A$ and $x' \in B^*$ such that $x'a = b'$. Then for all $b'' \in B^{**}$ we have

$$\langle\pi_\ell^{***}(e'', b''), b'\rangle = \langle e'', \pi_\ell^{**}(b'', b')\rangle = \lim_\alpha\langle\pi_\ell^{**}(b'', b'), e_\alpha\rangle$$
$$= \lim_\alpha\langle b'', \pi_\ell^{*}(b', e_\alpha)\rangle = \lim_\alpha\langle b'', \pi_\ell^{*}(x'a, e_\alpha)\rangle$$

$$= \lim_{\alpha}\langle b'', \pi_\ell^*(x', ae_\alpha)\rangle = \lim_{\alpha}\langle \pi_\ell^{**}(b'', x'), ae_\alpha\rangle$$
$$= \langle \pi_\ell^{**}(b'', x'), a\rangle = \langle b'', b'\rangle.$$

Thus $\pi_\ell^{***}(e'', b'') = b''$ consequently B^{**} has left unit $A^{**} - module$. It follows that $e'' \in Z_{B^{**}}(A^{**})$. If we take $Z_{B^{**}}(A^{**}) = Z^t_{B^{**}}(A^{**})$, then $e'' \in Z^t_{B^{**}}(A^{**})$. Then the mapping $b'' \to \pi_r^{t***t}(b'', e'')$ is $weak^* - to - weak^*$ continuous from B^{**} into B^{**}. Since $e_\alpha \overset{w^*}{\to} e''$, $\pi_r^{t***t}(b'', e_\alpha) \overset{w^*}{\to} \pi_r^{t***t}(b'', e'')$. Let $b' \in B^*$ and $(b_\beta)_\beta \subseteq B$ such that $b_\beta \overset{w^*}{\to} b''$. Since $Z^t_{e''}(B^{**}) = B^{**}$, we have the following quality

$$\langle \pi_r^{t***t}(b'', e''), b'\rangle = \lim_{\alpha}\langle \pi_r^{t***t}(b'', e_\alpha), b'\rangle = \lim_{\alpha}\langle \pi_r^{t***}(e_\alpha, b''), b'\rangle$$
$$= \lim_{\alpha}\lim_{\beta}\langle \pi_r^{t***}(e_\alpha, b_\beta), b'\rangle = \lim_{\alpha}\lim_{\beta}\langle \pi_r(b_\beta, e_\alpha), b'\rangle$$
$$= \lim_{\alpha}\lim_{\beta}\langle b', \pi_r(b_\beta, e_\alpha)\rangle = \lim_{\beta}\lim_{\alpha}\langle b', \pi_r(b_\beta, e_\alpha)\rangle$$
$$= \lim_{\beta}\langle b', b_\beta\rangle = \langle b'', b'\rangle.$$

Thus $\pi_r^{t***t}(b'', e'') = \pi_r^{***}(b'', e'') = b''$. It follows that B^{**} has a right unit. Suppose that $b'' \in B^{**}$ and $(b_\beta)_\beta \subseteq B$ such that $b_\beta \overset{w^*}{\to} b''$. Then for all $b' \in B^*$ we have

$$\langle b'', b'\rangle = \langle \pi_r^{***}(b'', e''), b'\rangle = \langle b'', \pi_r^{**}(e'', b')\rangle = \lim_{\beta}\langle \pi_r^{**}(e'', b'), b_\beta\rangle$$
$$= \lim_{\beta}\langle e'', \pi_r^*(b', b_\beta)\rangle = \lim_{\beta}\lim_{\alpha}\langle \pi_r^*(b', b_\beta), e_\alpha\rangle$$
$$= \lim_{\beta}\lim_{\alpha}\langle \pi_r^*(b', b_\beta), e_\alpha\rangle = \lim_{\beta}\lim_{\alpha}\langle b', \pi_r(b_\beta, e_\alpha)\rangle$$
$$= \lim_{\alpha}\lim_{\beta}\langle \pi_r^{***}(b_\beta, e_\alpha), b'\rangle = \lim_{\alpha}\lim_{\beta}\langle b_\beta, \pi_r^{**}(e_\alpha, b')\rangle$$
$$= \lim_{\alpha}\langle b'', \pi_r^{**}(e_\alpha, b')\rangle.$$

It follows that $weak - \lim_\alpha \pi_r^{**}(e_\alpha, b') = b'$. So by Cohen factorization theorem, B^* factors on the right that is contradiction. □

Corollary 2-5. Let B be a Banach $A - bimodule$ and $e'' \in A^{**}$ be a mixed unit for B^{**}. If B^* factors on the left, but not on the right, then $Z_{B^{**}}(A^{**}) \neq Z^t_{B^{**}}(A^{**})$.

For a Banach algebra A, if e'' is a left mixed unit for A^{**}, then it is clear that $Z^t_{e''}(A^{**}) = A^{**}$. Then by using preceding theorem, we obtain Proposition 2.10 from [14].

We say that a Banach space B is weakly complete, if for every $(b_\alpha)_\alpha \subseteq B$, we have $b_\alpha \overset{w}{\to} b''$ in B^{**}, it follows that $b'' \in B$.

Theorem 2-6. Suppose that B is a weakly complete Banach space. Then we have the following assertions.

1. Let B be a Banach left $A - module$ and B^{**} has a left mixed unit $e'' \in A^{**}$. If $AB^{**} \subseteq B$, then B is reflexive.

(2) Let B be a Banach right $A-module$ and B^{**} has a right mixed unit $e'' \in A^{**}$. If $Z_{A^{**}}(B^{**})A \subseteq B$, then $Z_{A^{**}}(B^{**}) = B$.

Proof. (1) Assume that $b'' \in B^{**}$. Since e'' is also mixed unit for A^{**}, there is a BAI $(e_\alpha)_\alpha \subseteq A$ for A such that $e_\alpha \overset{w^*}{\rightarrow} e''$. Then $\pi_\ell^{***}(e_\alpha, b'') \overset{w^*}{\rightarrow} \pi_\ell^{***}(e'', b'') = b''$ in B^{**}. Since $AB^{**} \subseteq B$, we have $\pi_\ell^{***}(e_\alpha, b'') \in B$. Consequently $\pi_\ell^{***}(e_\alpha, b'') \overset{w}{\rightarrow} \pi_\ell^{***}(e'', b'') = b''$ in B. Since B is a weakly complete, $b'' \in B$, and so B is reflexive.

(2) Since $b'' \in Z_{A^{**}}(B^{**})$, we have $\pi_r^{***}(b'', e_\alpha) \overset{w^*}{\rightarrow} \pi_r^{***}(b'', e'') = b''$ in B^{**}. Since $Z_{A^{**}}(B^{**})A \subseteq B$, $\pi_r^{***}(b'', e_\alpha) \in B$. Consequently we have $\pi_r^{***}(b'', e_\alpha) \overset{w}{\rightarrow} \pi_r^{***}(b'', e'') = b''$ in B. It follows that $b'' \in B$, since B is a weakly complete. □

A functional a' in A^* is said to be *wap* (weakly almost periodic) on A if the mapping $a \to a'a$ from A into A^* is weakly compact. The preceding definition is equivalent to the following condition, see [6, 14, 18].
For any two net $(a_\alpha)_\alpha$ and $(b_\beta)_\beta$ in $\{a \in A : \parallel a \parallel \leq 1\}$, we have

$$lim_\alpha lim_\beta \langle a', a_\alpha b_\beta \rangle = lim_\beta lim_\alpha \langle a', a_\alpha b_\beta \rangle,$$

whenever both iterated limits exist. The collection of all *wap* functionals on A is denoted by $wap(A)$. Also we have $a' \in wap(A)$ if and only if $\langle a''b'', a' \rangle = \langle a''ob'', a' \rangle$ for every a'', $b'' \in A^{**}$.

Definition 2-7. Let B be a Banach left $A-module$. Then, $b' \in B^*$ is said to be left weakly almost periodic functional if the set $\{\pi_\ell^*(b', a) : \ a \in A, \ \parallel a \parallel \leq 1\}$ is relatively weakly compact. We denote by $wap_\ell(B)$ the closed subspace of B^* consisting of all the left weakly almost periodic functionals in B^*.
The definition of the right weakly almost periodic functional ($= wap_r(B)$) is the same. By [18], the definition of $wap_\ell(B)$ is equivalent to the following

$$\langle \pi_\ell^{***}(a'', b''), b' \rangle = \langle \pi_\ell^{t***t}(a'', b''), b' \rangle$$

for all $a'' \in A^{**}$ and $b'' \in B^{**}$. Thus, we can write

$$wap_\ell(B) = \{b' \in B^* : \ \langle \pi_\ell^{***}(a'', b''), b' \rangle = \langle \pi_\ell^{t***t}(a'', b''), b' \rangle \\ for \ all \ a'' \in A^{**}, \ b'' \in B^{**}\}.$$

Theorem 2-8. Suppose that B is a Banach left $A-module$. Consider the following statements.

(1) $B^*A \subseteq wap_\ell(B)$.
(2) $AA^{**} \subseteq Z_{B^{**}}(A^{**})$.
(3) $AA^{**} \subseteq AZ_{B^{**}}((A^*A)^*)$.

Then, we have (1) $\Leftrightarrow$ (2) $\Leftarrow$ (3).

Proof. (1) $\Rightarrow$ (2)
Let $(b''_\alpha)_\alpha \subseteq B^{**}$ such that $b''_\alpha \overset{w^*}{\to} b''$. Then for all $a \in A$ and $a'' \in A^{**}$, we have

$$\langle \pi_\ell^{***}(aa'', b''_\alpha), b'\rangle = \langle aa'', \pi_\ell^{**}(b''_\alpha, b')\rangle = \langle a'', \pi_\ell^{**}(b''_\alpha, b')a\rangle$$
$$= \langle a'', \pi_\ell^{**}(b''_\alpha, b'a)\rangle = \langle \pi_\ell^{***}(a'', b''_\alpha), b'a\rangle \to \langle \pi_\ell^{***}(a'', b''), b'a\rangle$$
$$= \langle \pi_\ell^{***}(aa'', b''), b'\rangle.$$

Hence $aa'' \in Z_{B^{**}}(A^{**})$.
(2) $\Rightarrow$ (1)
Let $a \in A$ and $b' \in B^*$. Then

$$\langle \pi_\ell^{***}(a'', b''), b'a\rangle = \langle a\pi_\ell^{***}(a'', b''), b'\rangle = \langle \pi_\ell^{***}(aa'', b''), b'\rangle$$
$$= \langle \pi_\ell^{t***t}(aa'', b''), b'\rangle = \langle \pi_\ell^{t***t}(a'', b''), b'a\rangle.$$

It follow that $b'a \in wap_\ell(B)$.
(3) $\Rightarrow$ (2)
Since $AZ_{B^{**}}((A^*A)^*) \subseteq Z_{B^{**}}(A^{**})$, proof is hold. $\square$

In the preceding theorem, if we take $B = A$, then we obtain Theorem 3.6 from [14] and the same as preceding theorem, we can claim the following assertions:
If B is a Banach right $A - module$, then for the following statements we have (1) $\Leftrightarrow$ (2) $\Leftarrow$ (3).

1. $AB^* \subseteq wap_r(B)$.
2. $A^{**}A \subseteq Z_{B^{**}}(A^{**})$.
3. $A^{**}A \subseteq Z_{B^{**}}((A^*A)^*)A$.

The proof of the this assertion is similar to proof of Theorem 2-8.

Corollary 2-9. Suppose that B is a Banach $A - bimodule$. Then if A is a left [resp. right] ideal in A^{**}, then $B^*A \subseteq wap_\ell(B)$ [resp. $AB^* \subseteq wap_r(B)$].

Example 2-10. Suppose that $1 \leq p \leq \infty$ and q is conjugate of p. We know that if G is compact, then $L^1(G)$ is a two-sided ideal in its second dual of it. By preceding Theorem we have $L^q(G) * L^1(G) \subseteq wap_\ell(L^p(G))$ and $L^1(G) * L^q(G) \subseteq wap_r(L^p(G))$. Also if G is finite, then $L^q(G) \subseteq wap_\ell(L^p(G)) \cap wap_r(L^p(G))$. Hence we conclude that

$$Z_{L^1(G)^{**}}(L^p(G)^{**}) = L^p(G) \ and \ Z_{L^p(G)^{**}}(L^1(G)^{**}) = L^1(G).$$

Theorem 2-11. We have the following assertions.

1. Suppose that B is a Banach left $A - module$ and $b' \in B^*$. Then $b' \in wap_\ell(B)$ if and only if the adjoint of the mapping $\pi_\ell^*(b', \ ) : A \to B^*$ is $weak^*-to-weak$ continuous.
2. Suppose that B is a Banach right $A-module$ and $b' \in B^*$. Then $b' \in wap_r(B)$ if and only if the adjoint of the mapping $\pi_r^*(b', \ ) : B \to A^*$ is $weak^*-to-weak$ continuous.

Proof. (1) Assume that $b' \in wap_\ell(B)$ and $\pi_\ell^*(b',\ )^* : B^{**} \to A^*$ is the adjoint of $\pi_\ell^*(b',\ )$. Then for every $b'' \in B^{**}$ and $a \in A$, we have

$$\langle \pi_\ell^*(b',\ )^* b'', a\rangle = \langle b'', \pi_\ell^*(b', a)\rangle.$$

Suppose $(b''_\alpha)_\alpha \subseteq B^{**}$ such that $b''_\alpha \overset{w^*}{\to} b''$ and $a'' \in A^{**}$ and $(a_\beta)_\beta \subseteq A$ such that $a_\beta \overset{w^*}{\to} a''$. By easy calculation, for all $y'' \in B^{**}$ and $y' \in B^*$, we have

$$\langle \pi_\ell^*(y',\ )^*, y''\rangle = \pi_\ell^{**}(y'', y').$$

Since $b' \in wap_\ell(B)$,

$$\langle \pi_\ell^{***}(a'', b''_\alpha), b'\rangle \to \langle \pi_\ell^{***}(a'', b''), b'\rangle.$$

Then we have the following statements

$$\lim_\alpha \langle a'', \pi_\ell^*(b',\ )^* b''_\alpha\rangle = \lim_\alpha \langle a'', \pi_\ell^{**}(b''_\alpha, b')\rangle$$
$$= \lim_\alpha \langle \pi_\ell^{***}(a'', b''_\alpha), b'\rangle = \langle \pi_\ell^{***}(a'', b''), b'\rangle$$
$$= \langle a'', \pi_\ell^*(b',\ )^* b''\rangle.$$

It follow that the adjoint of the mapping $\pi_\ell^*(b',\ ) : A \to B^*$ is $weak^* - to - weak$ continuous.
Conversely, let the adjoint of the mapping $\pi_\ell^*(b',\ ) : A \to B^*$ is $weak^* - to - weak$ continuous. Suppose $(b''_\alpha)_\alpha \subseteq B^{**}$ such that $b''_\alpha \overset{w^*}{\to} b''$ and $b' \in B^*$. Then for every $a'' \in A^{**}$, we have

$$\lim_\alpha \langle \pi_\ell^{***}(a'', b''_\alpha), b'\rangle = \lim_\alpha \langle a'', \pi_\ell^{**}(b''_\alpha, b')\rangle$$
$$= \lim_\alpha \langle a'', \pi_\ell^*(b',\ )^* b''_\alpha\rangle = \langle a'', \pi_\ell^*(b',\ )^* b''\rangle = \langle \pi_\ell^{***}(a'', b''), b'\rangle.$$

It follow that $b' \in wap_\ell(B)$.

(2) proof is similar to (1).

□

Corollary 2-12. Let A be a Banach algebra. Assume that $a' \in A^*$ and $T_{a'}$ is the linear operator from A into A^* defined by $T_{a'}a = a'a$. Then, $a' \in wap(A)$ if and only if the adjoint of $T_{a'}$ is $weak^* - to - weak$ continuous. So A is Arens regular if and only if the adjoint of the mapping $T_{a'}a = a'a$ is $weak^* - to - weak$ continuous for every $a' \in A^*$.

3. Lw^*w-property and Rw^*w-property

In this section, we introduce the new definition as $Left - weak^* - to - weak$ property and $Right - weak^* - to - weak$ property for Banach algebra A and make some relations between these concepts and topological centers of module actions. As some conclusion, for locally compact group G, if $a \in M(G)$ has Lw^*w- property [resp. Rw^*w-

property], then we have $L^1(G)^{**} * a \neq L^1(G)^{**}$ [resp. $a * L^1(G)^{**} \neq L^1(G)^{**}$], and also we have $Z_{L^1(G)^{**}}(M(G)^{**}) \neq M(G)^{**}$ and $Z_{M(G)^{**}}(L^1(G)^{**}) = L^1(G)$. For a finite group G, we have $Z_{M(G)^{**}}(L^1(G)^{**}) = L^1(G)^{**}$ and $Z_{L^1(G)^{**}}(M(G)^{**}) = M(G)^{**}$.

Definition 3-1. Let B be a Banach left $A - module$. We say that $a \in A$ has $Left - weak^* - to - weak$ property ($= Lw^*w-$ property) with respect to B, if for all $(b'_\alpha)_\alpha \subseteq B^*$, $ab'_\alpha \overset{w^*}{\rightarrow} 0$ implies $ab'_\alpha \overset{w}{\rightarrow} 0$. If every $a \in A$ has Lw^*w- property with respect to B, then we say that A has Lw^*w- property with respect to B. The definition of the $Right - weak^* - to - weak$ property ($= Rw^*w-$ property) is the same.
We say that $a \in A$ has $weak^* - to - weak$ property ($= w^*w-$ property) with respect to B if it has Lw^*w- property and Rw^*w- property with respect to B.
If $a \in A$ has Lw^*w- property with respect to itself, then we say that $a \in A$ has Lw^*w- property.

For preceding definition, we have some examples and remarks as follows.
a) If B is a Banach A-bimodule and reflexive, then A has w^*w-property with respect to B. Then we have the following statenents for group algebras.
i) $L^1(G)$, $M(G)$ and $A(G)$ have w^*w-property when G is finite.
ii) Let G be locally compact group. $L^1(G)$ [resp. $M(G)$] has w^*w-property [resp. Lw^*w- property] with respect to $L^p(G)$ whenever $p > 1$.
b) Suppose that B is a Banach left $A - module$ and e is left unit element of A such that $eb = b$ for all $b \in B$. If e has Lw^*w- property, then B is reflexive.
c) If S is a compact semigroup, then $C^+(S) = \{f \in C(S) :\ f > 0\}$ has w^*w-property.

Theorem 3-2. Suppose that B is a Banach $A - bimodule$. Then we have the following assertions.

(1) If $A^{**} = a_0 A^{**}$ [resp. $A^{**} = A^{**} a_0$] for some $a_0 \in A$ and a_0 has Rw^*w- property [resp. Lw^*w- property], then $Z_{B^{**}}(A^{**}) = A^{**}$.
(2) If $B^{**} = a_0 B^{**}$ [resp. $B^{**} = B^{**} a_0$] for some $a_0 \in A$ and a_0 has Rw^*w- property [resp. Lw^*w- property] with respect to B, then $Z_{A^{**}}(B^{**}) = B^{**}$.

Proof. (1) Suppose that $A^{**} = a_0 A^{**}$ for some $a_0 \in A$ and a_0 has Rw^*w- property. Let $(b''_\alpha)_\alpha \subseteq B^{**}$ such that $b''_\alpha \overset{w^*}{\rightarrow} b''$. Then for all $a \in A$ and $b' \in B^*$, we have

$$\langle \pi_\ell^{**}(b''_\alpha, b'), a\rangle = \langle b''_\alpha, \pi_\ell^*(b', a)\rangle \rightarrow \langle b'', \pi_\ell^*(b', a)\rangle = \langle \pi_\ell^{**}(b'', b'), a\rangle,$$

it follow that $\pi_\ell^{**}(b''_\alpha, b') \overset{w^*}{\rightarrow} \pi_\ell^{**}(b'', b')$. Also we can write $\pi_\ell^{**}(b''_\alpha, b')a_0 \overset{w^*}{\rightarrow} \pi_\ell^{**}(b'', b')a_0$. Since a_0 has Rw^*w- property, $\pi_\ell^{**}(b''_\alpha, b')a_0 \overset{w}{\rightarrow} \pi_\ell^{**}(b'', b')a_0$. Now let $a'' \in A^{**}$. Then there is $x'' \in A^{**}$ such that $a'' = a_0 x''$ consequently we have

$$\begin{aligned}\langle \pi_\ell^{***}(a'', b''_\alpha), b'\rangle &= \langle a'', \pi_\ell^{**}(b''_\alpha, b')\rangle = \langle x'', \pi_\ell^{**}(b''_\alpha, b')a_0\rangle \\ &\rightarrow \langle x'', \pi_\ell^{**}(b'', b')a_0\rangle = \langle \pi_\ell^{***}(a'', b''_\alpha), b'\rangle.\end{aligned}$$

We conclude that $a'' \in Z_{B^{**}}(A^{**})$. Proof of the next part is the same as the preceding proof.

(2) Let $B^{**} = a_0 B^{**}$ for some $a_0 \in A$ and a_0 has Rw^*w- property with respect to B. Assume that $(a''_\alpha)_\alpha \subseteq A^{**}$ such that $a''_\alpha \overset{w^*}{\to} a''$. Then for all $b \in B$, we have

$$\langle \pi_r^{**}(a''_\alpha, b'), b\rangle = \langle a''_\alpha, \pi_r^*(b', b)\rangle \to \langle a'', \pi_r^*(b', b)\rangle = \langle \pi_r^{**}(a'', b'), b\rangle.$$

We conclude that $\pi_r^{**}(a''_\alpha, b') \overset{w^*}{\to} \pi_r^{**}(a'', b')$ then we have $\pi_r^{**}(a''_\alpha, b')a_0 \overset{w^*}{\to} \pi_r^{**}(a'', b')a_0$. Since a_0 has Rw^*w- property with respect to B, $\pi_r^{**}(a''_\alpha, b')a_0 \overset{w}{\to} \pi_r^{**}(a'', b')a_0$.
Now let $b'' \in B^{**}$. Then there is $x'' \in B^{**}$ such that $b'' = a_0 x''$. Hence, we have

$$\begin{aligned} \langle \pi_r^{***}(b'', a''_\alpha), b'\rangle &= \langle b'', \pi_r^{**}(a''_\alpha, b')\rangle = \langle a_0 x'', \pi_r^{**}(a''_\alpha, b')\rangle \\ = \langle x'', \pi_r^{**}(a''_\alpha, b')a_0\rangle &\to \langle x'', \pi_r^{**}(a'', b')a_0\rangle = \langle b'', \pi_r^{**}(a'', b')\rangle \\ &= \langle \pi_r^{***}(b'', a''), b'\rangle. \end{aligned}$$

It follow that $b'' \in Z_{A^{**}}(B^{**})$. The next part is similar to the preceding proof.

□

Example 3-3.

(1) By using Theorem 2-3, for locally compact group G, if $a \in M(G)$ has Lw^*w- property [resp. Rw^*w- property], then we have $L^1(G)^{**} * a \neq L^1(G)^{**}$ [resp. $a * L^1(G)^{**} \neq L^1(G)^{**}$].

(2) If G is finite, then by Theorem 2-3, we have $Z_{M(G)^{**}}(L^1(G)^{**}) = L^1(G)^{**}$ and $Z_{L^1(G)^{**}}(M(G)^{**}) = M(G)^{**}$.

Theorem 3-4. Suppose that B is a Banach $A-$ *bimodule* and A has a BAI. Then we have the following assertions.

(1) If B^* factors on the left [resp. right] with respect to A and A has Rw^*w- property [resp. Lw^*w- property], then $Z_{B^{**}}(A^{**}) = A^{**}$.

(2) If B^* factors on the left [resp. right] with respect to A and A has Rw^*w- property [resp. Lw^*w- property] with respect B, then $Z_{A^{**}}(B^{**}) = B^{**}$.

Proof. (1) Assume that B^* factors on the left and A has Rw^*w- property. Let $(b''_\alpha)_\alpha \subseteq B^{**}$ such that $b''_\alpha \overset{w^*}{\to} b''$. Since $B^*A = B^*$, for all $b' \in B^*$ there are $x \in A$ and $y' \in B^*$ such that $b' = y'x$. Then for all $a \in A$, we have

$$\begin{aligned} &\langle \pi_\ell^{**}(b''_\alpha, y')x, a\rangle = \langle b''_\alpha, \pi_\ell^*(y', a)x\rangle = \langle \pi_\ell^{**}(b''_\alpha, b'), a\rangle \\ &= \langle b''_\alpha, \pi_\ell^*(b', a)\rangle \to \langle b'', \pi_\ell^*(b', a)\rangle = \langle \pi_\ell^{**}(b'', y')x, a\rangle. \end{aligned}$$

Thus, we conclude that $\pi_\ell^{**}(b''_\alpha, y')x \overset{w^*}{\to} \pi_\ell^{**}(b'', y')x$. Since A has Rw^*w- property, $\pi_\ell^{**}(b''_\alpha, y')x \overset{w}{\to} \langle \pi_\ell^{**}(b'', y')x$. Now let $a'' \in A^{**}$. Then

$$\langle \pi_\ell^{***}(a'', b''_\alpha), b'\rangle = \langle a'', \pi_\ell^{**}(b''_\alpha, b')\rangle = \langle a'', \pi_\ell^{**}(b''_\alpha, y')x\rangle$$

$$\rightarrow \langle a'', \pi_\ell^{**}(b'', y')x\rangle = \langle \pi_\ell^{***}(a'', b''), b'\rangle.$$

It follow that $a'' \in Z_{B^{**}}(A^{**})$.
If B^* factors on the right with respect to A, and assume that A has Lw^*w- property, then proof is the same as preceding proof.

(2) Let B^* factors on the left with respect to A and A has Rw^*w- property with respect to B. Assume that $(a''_\alpha)_\alpha \subseteq A^{**}$ such that $a''_\alpha \overset{w^*}{\rightarrow} a''$. Since $B^*A = B^*$, for all $b' \in B^*$ there are $x \in A$ and $y' \in B^*$ such that $b' = y'x$. Then for all $b \in B$, we have

$$\begin{aligned}\langle \pi_r^{**}(a''_\alpha, y')x, b\rangle &= \langle \pi_r^{**}(a''_\alpha, b'), b\rangle = \langle a''_\alpha, \pi_r^*(b', b)\rangle \\ &= \langle a'', \pi_r^*(b', b)\rangle = \langle \pi_r^{**}(a'', y')x, b\rangle.\end{aligned}$$

Consequently $\pi_r^{**}(a''_\alpha, y')x \overset{w^*}{\rightarrow} \pi_r^{**}(a'', y')x$. Since A has Rw^*w- property with respect to B, $\pi_r^{**}(a''_\alpha, y')x \overset{w}{\rightarrow} \pi_r^{**}(a'', y')x$. It follow that for all $b'' \in B^{**}$, we have

$$\begin{aligned}\langle \pi_r^{***}(b'', a''_\alpha), b'\rangle = \langle b'', \pi_r^{**}(a''_\alpha, y')x\rangle &\rightarrow \langle b'', \pi_r^{**}(a'', y')x\rangle \\ &= \langle \pi_r^{***}(b'', a''), b'\rangle.\end{aligned}$$

Thus we conclude that $b'' \in Z_{A^{**}}(B^{**})$.
The proof of the next assertions is the same as the preceding proof.

□

Theorem 3-5. Suppose that B is a Banach $A-$*bimodule*. Then we have the following assertions.

(1) If $a_0 \in A$ has Rw^*w- property with respect to B, then $a_0A^{**} \subseteq Z_{B^{**}}(A^{**})$ and $a_0B^* \subseteq wap_\ell(B)$.
(2) If $a_0 \in A$ has Lw^*w- property with respect to B, then $A^{**}a_0 \subseteq Z_{B^{**}}(A^{**})$ and $B^*a_0 \subseteq wap_\ell(B)$.
(3) If $a_0 \in A$ has Rw^*w- property with respect to B, then $a_0B^{**} \subseteq Z_{A^{**}}(B^{**})$ and $B^*a_0 \subseteq wap_r(B)$.
(4) If $a_0 \in A$ has Lw^*w- property with respect to B, then $B^{**}a_0 \subseteq Z_{A^{**}}(B^{**})$ and $a_0B^* \subseteq wap_r(B)$.

Proof. (1) Let $(b''_\alpha)_\alpha \subseteq B^{**}$ such that $b''_\alpha \overset{w^*}{\rightarrow} b''$. Then for all $a \in A$ and $b' \in B^*$, we have

$$\begin{aligned}\langle \pi_\ell^{**}(b''_\alpha, b')a_0, a\rangle &= \langle \pi_\ell^{**}(b''_\alpha, b'), a_0a\rangle = \langle b''_\alpha, \pi_\ell^*(b', a_0a)\rangle \\ &\rightarrow \langle b'', \pi_\ell^*(b', a_0a)\rangle = \langle \pi_\ell^{**}(b'', b')a_0, a\rangle.\end{aligned}$$

It follow that $\pi_\ell^{**}(b''_\alpha, b')a_0 \overset{w^*}{\rightarrow} \pi_\ell^{**}(b'', b')a_0$. Since a_0 has Rw^*w- property with respect to B, $\pi_\ell^{**}(b''_\alpha, b')a_0 \overset{w}{\rightarrow} \pi_\ell^{**}(b'', b')a_0$.
We conclude that $a_0a'' \in Z_{B^{**}}(A^{**})$ so that $a_0A^{**} \in Z_{B^{**}}(A^{**})$. Since $\pi_\ell^{**}(b'', b')a_0 = \pi_\ell^{**}(b'', b'a_0)$, $a_0B^* \subseteq wap_\ell(B)$.

(2) proof is similar to (1).

(3) Assume that $(a''_\alpha)_\alpha\subseteq A^{**}$ such that $a''_\alpha\overset{w^*}{\to}a''$. Let $b\in B$ and $b'\in B^*$. Then we have

$$\begin{aligned}\langle\pi_r^{**}(a''_\alpha,b')a_0,b\rangle&=\langle\pi_r^{**}(a''_\alpha,b'),a_0b\rangle=\langle a''_\alpha,\pi_r^{*}(b',a_0b)\rangle\\&\to\langle a'',\pi_r^{*}(b',a_0b)\rangle=\langle\pi_r^{**}(a'',b')a_0,b\rangle.\end{aligned}$$

Thus we conclude $\pi_r^{**}(a''_\alpha,b')a_0\overset{w^*}{\to}\pi_r^{**}(a'',b')a_0$. Since a_0 has Rw^*w- property with respect to B, $\pi_r^{**}(a''_\alpha,b')a_0\overset{w}{\to}\pi_r^{**}(a'',b')a_0$. If $b''\in B^{**}$, then we have

$$\begin{aligned}\langle\pi_r^{***}(a_0b'',a''_\alpha),b'\rangle&=\langle a_0b'',\pi_r^{***}(a''_\alpha,b')\rangle=\langle b'',\pi_r^{**}(a''_\alpha,b')a_0\rangle\\&=\langle b'',\pi_r^{**}(a''_\alpha,b')a_0\rangle=\langle\pi_r^{***}(a_0b'',a''),b'\rangle.\end{aligned}$$

It follow that $a_0b''\in Z_{A^{**}}(B^{**})$. Consequently we have $a_0B^{**}\in Z_{A^{**}}(B^{**})$. The proof of the next assertion is clear.

(4) Proof is similar to (3).

□

Theorem 3-6. Let B be a Banach $A-$ *bimodule.* Then we have the following assertions.

(1) Suppose

$$\lim_\alpha\lim_\beta\langle b'_\beta,b_\alpha\rangle=\lim_\beta\lim_\alpha\langle b'_\beta,b_\alpha\rangle,$$

for every $(b_\alpha)_\alpha\subseteq B$ and $(b'_\beta)_\beta\subseteq B^*$. Then A has Lw^*w- property and Rw^*w- property with respect to B.

(2) If for some $a\in A$,

$$\lim_\alpha\lim_\beta\langle ab'_\beta,b_\alpha\rangle=\lim_\beta\lim_\alpha\langle ab'_\beta,b_\alpha\rangle,$$

for every $(b_\alpha)_\alpha\subseteq B$ and $(b'_\beta)_\beta\subseteq B^*$, then a has Rw^*w- property with respect to B. Also if for some $a\in A$,

$$\lim_\alpha\lim_\beta\langle b'_\beta a,b_\alpha\rangle=\lim_\beta\lim_\alpha\langle b'_\beta a,b_\alpha\rangle,$$

for every $(b_\alpha)_\alpha\subseteq B$ and $(b'_\beta)_\beta\subseteq B^*$, then a has Lw^*w- property with respect to B.

Proof. (1) Assume that $a\in A$ such that $ab'_\beta\overset{w^*}{\to}0$ where $(b'_\beta)_\beta\subseteq B^*$. Let $b''\in B^{**}$ and $(b_\alpha)_\alpha\subseteq B$ such that $b_\alpha\overset{w^*}{\to}b''$. Then

$$\begin{aligned}\lim_\beta\langle b'',ab'_\beta\rangle&=\lim_\beta\lim_\alpha\langle b_\alpha,ab'_\beta\rangle=\lim_\beta\lim_\alpha\langle ab'_\beta,b_\alpha\rangle\\&=\lim_\alpha\lim_\beta\langle ab'_\beta,b_\alpha\rangle=0.\end{aligned}$$

We conclude that $ab'_\beta\overset{w}{\to}0$, so A has Lw^*w- property. It also easy that A has Rw^*w- property.

(2) Proof is easy and is the same as (1).

□

Definition 3-7. Let B be a Banach left $A-module$. We say that B^* strong factors on the left [resp. right] if for all $(b'_\alpha)_\alpha \subseteq B^*$ there are $(a_\alpha)_\alpha \subseteq A$ and $b' \in B^*$ such that $b'_\alpha = b'a_\alpha$ [resp. $b'_\alpha = a_\alpha b'$] where $(a_\alpha)_\alpha$ has limit the $weak^*$ topology in A^{**}.
If B^* strong factors on the left and right, then we say that B^* strong factors on the both side.
It is clear that if B^* strong factors on the left [resp. right], then B^* factors on the left [resp. right].

Theorem 3-8. Suppose that B is a Banach $A-bimodule$. Assume that $AB^* \subseteq wap_\ell B$. If B^* strong factors on the left [resp. right], then A has Lw^*w- property [resp. Rw^*w- property] with respect to B.

Proof. Let $(b'_\alpha)_\alpha \subseteq B^*$ such that $ab'_\alpha \overset{w^*}{\rightarrow} 0$. Since B^* strong factors on the left, there are $(a_\alpha)_\alpha \subseteq A$ and $b' \in B^*$ such that $b'_\alpha = b'a_\alpha$. Let $b'' \in B^{**}$ and $(b_\beta)_\beta \subseteq B$ such that $b_\beta \overset{w^*}{\rightarrow} b''$. Then we have

$$\begin{aligned}\lim_\alpha \langle b'', ab'_\alpha\rangle &= \lim_\alpha \lim_\beta \langle b_\beta, ab'_\alpha\rangle = \lim_\alpha \lim_\beta \langle ab'_\alpha, b_\beta\rangle \\ &= \lim_\alpha \lim_\beta \langle ab'a_\alpha, b_\beta\rangle = \lim_\alpha \lim_\beta \langle ab', a_\alpha b_\beta\rangle \\ &= \lim_\beta \lim_\alpha \langle ab', a_\alpha b_\beta\rangle = \lim_\beta \lim_\alpha \langle ab'_\alpha, b_\beta\rangle = 0\end{aligned}$$

It follow that $ab'_\alpha \overset{w}{\rightarrow} 0$. □

Problems .

(1) Suppose that B is a Banach $A-bimodule$. If B is left or right factors with respect to A, dose A has Lw^*w-property or Rw^*w-property, respectively?
(2) Suppose that B is a Banach $A-bimodule$. Let A has Lw^*w-property with respect to B. Dose $Z_{B^{**}}(A^{**}) = A^{**}$?

4. Factorization properties and topological centers of left module actions

Let B be a left Banach $A-module$ and e be a left unit element of A. We say that e is a left unit (resp. weakly left unit) $A-module$ for B, if $\pi_\ell(e,b) = b$ (resp. $\langle b', \pi_\ell(e,b)\rangle = \langle b', b\rangle$ for all $b' \in B^*$) where $b \in B$. The definition of right unit (resp. weakly right unit) $A-module$ is similar.
We say that a Banach $A-bimodule$ B is a unital $A-module$, if B has left and right unit $A-module$ that are equal then we say that B is unital $A-module$.

Let B be a left Banach $A-module$ and $(e_\alpha)_\alpha \subseteq A$ be a LAI [resp. weakly left approximate identity(=WLAI)] for A. We say that $(e_\alpha)_\alpha$ is left approximate identity $(= LAI)$[resp. weakly left approximate identity $(=WLAI)$] for B, if for all $b \in B$, $\pi_\ell(e_\alpha, b) \to b$ (resp. $\pi_\ell(e_\alpha, b) \overset{w}{\to} b$). The definition of the right approximate identity $(= RAI)$[resp. weakly right approximate identity $(= WRAI)$] is similar.
$(e_\alpha)_\alpha \subseteq A$ is called a approximate identity $(= AI)$[resp. weakly approximate identity (WAI)] for B, if B has the same left and right approximate identity [resp. weakly left and right approximate identity].
Let $(e_\alpha)_\alpha \subseteq A$ be $weak^*$ left approximate identity for A^{**}. We say that $(e_\alpha)_\alpha$ is $weak^*$ left approximate identity as $A^{**}-module$ $(= W^*LAI\ as\ A^{**}-module)$ for B^{**}, if for all $b'' \in B^{**}$, we have $\pi_\ell^{***}(e_\alpha, b'') \overset{w^*}{\to} b''$. The definition of the $weak^*$ right approximate identity $(= W^*RAI\ )$ is similar.
$(e_\alpha)_\alpha \subseteq A$ is called a $weak^*$ approximate identity $(= W^*AI\ )$ for B^{**}, if B^{**} has the same $weak^*$ left and right approximate identity.
Let B be a Banach $A-bimodule$. We say that B is a left [resp. right] factors with respect to A, if $BA = B$ [resp. $AB = B$].

Theorem 4-1. Let B be a Banach left $A-module$ and $(e_\alpha)_\alpha \subseteq A$ be a $LBAI$ for B. Then the following assertions hold.

(1) For each $b' \in B^*$, we have $\pi_\ell^*(b', e_\alpha) \overset{w^*}{\to} b'$.
(2) B^* factors on the left with respect to A if and only if B^{**} has a W^*LBAI $(e_\alpha)_\alpha \subseteq A$.
(3) B^{**} has a W^*LBAI $(e_\alpha)_\alpha \subseteq A$ if and only if B^{**} has a left unit element $e'' \in A^{**}$ such that $e_\alpha \overset{w^*}{\to} e''$.

Proof. (1) Let $b \in B$ and $b' \in B^*$. Since $\mid \langle b', \pi_\ell(e_\alpha, b) \mid \leq \parallel b' \parallel \parallel \pi_\ell(e_\alpha, b) \parallel$, we have the following equality

$$\lim_\alpha \langle \pi_\ell^*(b', e_\alpha), b\rangle = \lim_\alpha \langle b', \pi_\ell(e_\alpha, b)\rangle = 0.$$

It follows that $\pi_\ell^*(b', e_\alpha) \overset{w^*}{\to} 0$.

(2) Let B^* factors on the left with respect to A. Then for every $b' \in B^*$, there are $x' \in B^*$ and $a \in A$ such that $b' = x'a$. Then for every $b'' \in B^{**}$, we have

$$\begin{aligned}\langle \pi_\ell^{***}(e_\alpha, b''), b'\rangle &= \langle e_\alpha, \pi_\ell^{**}(b'', b')\rangle = \langle \pi_\ell^{**}(b'', b'), e_\alpha\rangle \\ = \langle b'', \pi_\ell^*(b', e_\alpha)\rangle &= \langle b'', \pi_\ell^*(x'a, e_\alpha)\rangle = \langle b'', \pi_\ell^*(x', ae_\alpha)\rangle \\ &= \langle \pi_\ell^{**}(b'', x'), ae_\alpha\rangle \to \langle \pi_\ell^{**}(b'', x'), a\rangle \\ &= \langle b'', b'\rangle.\end{aligned}$$

It follows that

$$\pi_\ell^{***}(e_\alpha, b'') \overset{w^*}{\to} b'',$$

and so B^{**} has W^*LBAI.
Conversely, let $b' \in B^*$. Then for every $b'' \in B^{**}$, we have

$$\langle b'', \pi_\ell^*(b', e_\alpha)\rangle = \langle \pi_\ell^{***}(e_\alpha, b''), b'\rangle \to \langle b'', b'\rangle.$$

It follows that

$$\pi_\ell^*(b', e_\alpha) \xrightarrow{w} b',$$

and so by Cohen factorization theorem, we are done.

(3) Assume that B^{**} has a W^*LBAI $(e_\alpha)_\alpha \subseteq A$. Without loss generality, let $e'' \in A^{**}$ be a left unit for A^{**} with respect to the first Arens product such that $e_\alpha \xrightarrow{w^*} e''$. Then for each $b' \in B^*$, we have

$$\langle \pi_\ell^{***}(e'', b''), b'\rangle = \langle e'', \pi_\ell^{**}(b'', b')\rangle$$

$$= \lim_\alpha \langle e_\alpha, \pi_\ell^{**}(b'', b')\rangle = \lim_\alpha \langle \pi_\ell^{**}(b'', b'), e_\alpha\rangle$$

$$= \lim_\alpha \langle b'', \pi_\ell^{*}(b', e_\alpha)\rangle = \lim_\alpha \langle \pi_\ell^{****}(b', e_\alpha), b''\rangle$$

$$= \lim_\alpha \langle b', \pi_\ell^{***}(e_\alpha, b'')\rangle = \lim_\alpha \langle \pi_\ell^{***}(e_\alpha, b''), b'\rangle$$

$$= \langle b'', b'\rangle.$$

Thus $e'' \in A^{**}$ is a left unit for B^{**}.

Conversely, let $e'' \in A^{**}$ be a left unit for B^{**} and assume that $e_\alpha \xrightarrow{w^*} e''$ in A^{**}. Then for every $b'' \in B^{**}$ and $b' \in B^*$, we have

$$\langle \pi_\ell^{***}(e_\alpha, b''), b'\rangle = \langle e_\alpha, \pi_\ell^{**}(b'', b')\rangle$$

$$\to \langle e'', \pi_\ell^{**}(b'', b')\rangle = \langle \pi_\ell^{***}(e'', b''), b'\rangle$$

$$= \langle b'', b'\rangle.$$

It follows that $\pi_\ell^{***}(e_\alpha, b'') \xrightarrow{w^*} b''$.

□

Corollary 4-2. Let B be a Banach left $A-module$ and A has a $BLAI$. If B^{**} has a W^*LBAI , then

$$\{a'' \in A^{**} : \ Aa'' \subseteq A\} \subseteq Z_{B^{**}}(A^{**}).$$

Proof. By using the preceding theorem, since B^{**} has W^*LBAI, B^* factors on the left with respect to A. Suppose that $b' \in B^*$. Then there are $x' \in B^*$ and $a \in A$ such that $b' = x'a$. Assume that $a'' \in A^{**}$ such that $Aa'' \subseteq A$. Let $b'' \in B^{**}$ and $(b''_\alpha)_\alpha \subseteq B^{**}$ such that $b''_\alpha \xrightarrow{w^*} b''$ in B^{**}. Then we have the following equality

$$\lim_\alpha \langle \pi_\ell^{***}(a'', b''_\alpha), b'\rangle = \lim_\alpha \langle \pi_\ell^{***}(a'', b''_\alpha), x'a\rangle$$

$$= \lim_\alpha \langle a\pi_\ell^{***}(a'', b''_\alpha), x'\rangle = \langle \pi_\ell^{***}(aa'', b''), x'\rangle = \langle \pi_\ell^{***}(a'', b''), b'\rangle.$$

It follows that $a'' \in Z_{B^{**}}(A^{**})$. □

In the preceding corollary, if we take $B = A$, then we have the following conclusion

$$\{a'' \in A^{**} : \ Aa'' \subseteq A\} \subseteq Z_1(A^{**}).$$

Theorem 4-3. Let B be a Banach left $A-module$ and suppose that $b' \in wap_\ell(B)$. Let $a'' \in A^{**}$ and $(a_\alpha)_\alpha \subseteq A$ such that $a_\alpha \overset{w^*}{\to} a''$ in A^{**}. Then we have

$$\pi_\ell^*(b', a_\alpha) \overset{w}{\to} \pi_\ell^{****}(b', a'').$$

Proof. Assume that $b'' \in B^{**}$. Then we have the following equality

$$\langle \pi_\ell^{****}(b', a''), b''\rangle = \langle \pi_\ell^{***}(a'', b''), b'\rangle = \lim_\alpha \langle \pi_\ell^{***}(a_\alpha, b''), b'\rangle$$
$$= \lim_\alpha \langle b'', \pi_\ell^*(b', a_\alpha)\rangle.$$

Now suppose that $(b''_\beta)_\beta \subseteq B^{**}$ such that $b''_\beta \overset{w^*}{\to} b''$. Since $b' \in wap_\ell(B)$, we have

$$\langle \pi_\ell^{****}(b', a''), b''_\beta\rangle = \langle \pi_\ell^{***}(a'', b''_\beta), b'\rangle \to \langle \pi_\ell^{***}(a'', b''), b'\rangle$$
$$= \langle \pi_\ell^{****}(b', a''), b''\rangle.$$

Thus $\pi_\ell^{****}(b', a'') \in (B^{**}, weak^*)^* = B^*$. So we conclude that

$$\pi_\ell^*(b', a_\alpha) \overset{w}{\to} \pi_\ell^{****}(b', a'') \ in \ B^{**}.$$

□

In the preceding corollary, if we take $B = A$, then we obtain the following result. Suppose that $a' \in wap(A)$ and $a'' \in A^{**}$ such that $a_\alpha \overset{w^*}{\to} a''$ where $(a_\alpha)_\alpha \subseteq A$. Then we have $a'a_\alpha \overset{w}{\to} a'a''$.

Theorem 4-4. Let B be a Banach left $A-module$ and it has a $BLAI$ $(e_\alpha)_\alpha \subseteq A$. Suppose that $b' \in wap_\ell(B)$. Then we have

$$\pi_\ell^*(b', e_\alpha) \overset{w}{\to} b'.$$

Proof. Let $b'' \in B^{**}$ and $(b_\beta)_\beta \subseteq B$ such that $b_\beta \overset{w^*}{\to} b''$ in B^{**}. Then for every $b' \in wap_\ell(B)$, we have the following equality

$$\lim_\alpha \langle b'', \pi_\ell^*(b', e_\alpha)\rangle = \lim_\alpha \langle \pi_\ell^{****}(b', e_\alpha), b''\rangle$$
$$= \lim_\alpha \langle b', \pi_\ell^{***}(e_\alpha, b'')\rangle = \lim_\alpha \langle \pi_\ell^{***}(e_\alpha, b''), b'\rangle$$
$$= \lim_\alpha \lim_\beta \langle \pi_\ell(e_\alpha, b_\beta), b'\rangle = \lim_\beta \lim_\alpha \langle \pi_\ell(e_\alpha, b_\beta), b'\rangle$$
$$= \lim_\beta \langle b_\beta, b'\rangle = \langle b'', b'\rangle.$$

It follows that

$$\pi_\ell^*(b', e_\alpha) \overset{w}{\to} b'.$$

□

Corollary 4-5. Let B be a Banach left $A-module$ and it has a $BLAI$ $(e_\alpha)_\alpha \subseteq A$. Suppose that $wap_\ell(B) = B^*$. Then B^* factors on the left with respect to A.

Corollary 4-6. Let A be an Arens regular Banach algebra with $LBAI$. Then A^* factors on the left.

Example 4-7. i) Let G be finite group. Then we have the following equality

$$M(G)^*L^1(G) = M(G)^* \ and \ L^\infty(G)L^1(G) = L^\infty(G).$$

ii) Consider the Banach algebra $(\ell^1, .)$ that is Arens regular Banach algebra with unit element. Then we have $\ell^\infty.\ell^1 = \ell^\infty$.
iii) Let $\mathcal{K}(E)$ be a compact operators from E into E. Let $E = \ell^p$ with $p \in (1, \infty)$. Then by using [6, Theorem 2.6.23] and [19 , Example 2.3.18] , $\mathcal{K}(\ell^p)$ is Arens regular and amenable, respectively. Since $\mathcal{K}(\ell^p)$ is amenable, by using [19 , Proposition 2.2.1], $\mathcal{K}(\ell^p)$ has a BAI. Therefore by preceding corollary, $\mathcal{K}(\ell^p)^*$ factors on the left.

Theorem 4-8. Let B be a Banach left $A-module$ and B^* factors on the left with respect to A. If $AA^{**} \subseteq Z_{B^{**}}(A^{**})$, then $Z_{B^{**}}(A^{**}) = A^{**}$.

Proof. Let $b'' \in B^{**}$ and $(b''_\alpha)_\alpha \subseteq B^{**}$ such that $b''_\alpha \overset{w^*}{\to} b''$ in B^{**}. Suppose that $a'' \in A^{**}$. Since B^* factors on the left with respect to A, for every $b' \in B^*$, there are $x' \in B^*$ and $a \in A$ such that $b' = x'a$. Since $aa'' \in Z_{B^{**}}(A^{**})$, we have

$$\langle \pi_\ell^{***}(a'', b''_\alpha), b' \rangle = \langle \pi_\ell^{***}(a'', b''_\alpha), x'a \rangle$$
$$= \langle a\pi_\ell^{***}(a'', b''_\alpha), x' \rangle = \langle \pi_\ell^{***}(aa'', b''_\alpha), x' \rangle$$
$$\to \langle \pi_\ell^{***}(aa'', b''), x' \rangle = \langle \pi_\ell^{***}(a'', b''), b' \rangle.$$

It follows that

$$\pi_\ell^{***}(a'', b''_\alpha) \overset{w^*}{\to} \langle \pi_\ell^{***}(a'', b''),$$

and so $a'' \in Z_{B^{**}}(A^{**})$. □

Corollary 4-9. Let A be a Banach algebra and A^* factors on the left. If $AA^{**} \subseteq Z_1(A^{**})$, then A is Arens regular.

Theorem 4-10. Let B be a Banach left $A-module$ and B^{**} has a $LBAI$ with respect to A^{**}. Then B^{**} has a left unit with respect to A^{**}.

Proof. Assume that $(e''_\alpha)_\alpha \subseteq A^{**}$ is a $LBAI$ for B^{**}. By passing to a subnet, we may suppose that there is $e'' \in A^{**}$ such that $e''_\alpha \overset{w^*}{\to} e''$ in A^{**}. Then for every $b'' \in B^{**}$ and $b' \in B^*$, we have

$$\langle \pi_\ell^{***}(e'', b''), b' \rangle = \langle e'', \pi_\ell^{**}(b'', b') \rangle = \lim_\alpha \langle e''_\alpha, \pi_\ell^{**}(b'', b') \rangle$$

$$= \lim_{\alpha} \langle \pi_\ell^{***}(e''_\alpha, b''), b' \rangle = \langle b'', b' \rangle.$$

It follows that $\pi_\ell^{***}(e'', b'') = b''$. □

Corollary 4-11. Let A be a Banach algebra and A^{**} has a $LBAI$. Then A^{**} has a left unit with respect to the first Arens product.

Theorem 4-12. Let B be a Banach left $A - module$ and it has a $LBAI$ with respect to A. Then we have the following assertions.

1. B^* factors on the left with respect to A if and only if for each $b' \in B^*$, we have $\pi_\ell^*(b', e_\alpha) \overset{w}{\rightarrow} b'$ in B^*.
2. B factors on the left with respect to A if and only if for each $b \in B$, we have $\pi_\ell^*(b, e_\alpha) \overset{w}{\rightarrow} b$ in B.

Proof. (1) Assume that B^* factors on the left with respect to A. Then for every $b' \in B^*$, there are $x' \in B^*$ and $a \in A$ such that $b' = x'a$. Then for every $b'' \in B^{**}$, we have

$$\langle b'', \pi_\ell^*(b', e_\alpha) \rangle = \langle b'', \pi_\ell^*(x'a, e_\alpha) \rangle = \langle b'', \pi_\ell^*(x', ae_\alpha) \rangle$$

$$= \langle \pi_\ell^{**}(b'', x'), ae_\alpha) \rangle \rightarrow \langle \pi_\ell^{**}(b'', x'), a \rangle$$

$$= \langle b'', b' \rangle.$$

It follows that $\pi_\ell^*(b', e_\alpha) \overset{w}{\rightarrow} b'$ in B^*. The converse by Cohen factorization theorem hold.

(2) It is similar to the preceding proof.

□

In the preceding theorem, if we take $B = A$, we obtain Lemma 2.1 from [14].

Theorem 4-13. Let B be a Banach left $A - module$ and A has a $LBAI$ $(e_\alpha)_\alpha \subseteq A$ such that $e_\alpha \overset{w^*}{\rightarrow} e''$ in A^{**} where e'' is a left unit for A^{**}. Suppose that $Z^t_{e''}(B^{**}) = B^{**}$. Then, B factors on the right with respect to A if and only if e'' is a left unit for B^{**}.

Proof. Assume that B factors on the right with respect to A. Then for every $b \in B$, there are $x \in B$ and $a \in A$ such that $b = ax$. Then for every $b' \in B^*$, we have

$$\langle \pi_\ell^*(b', e_\alpha), b \rangle = \langle b', \pi_\ell(e_\alpha, b) \rangle = \langle \pi_\ell^{***}(e_\alpha, b), b' \rangle$$

$$= \langle \pi_\ell^{***}(e_\alpha, ax), b' \rangle = \langle \pi_\ell^{***}(e_\alpha a, x), b' \rangle$$

$$= \langle e_\alpha a, \pi_\ell^{**}(x, b') \rangle = \langle \pi_\ell^{**}(x, b'), e_\alpha a \rangle$$

$$\rightarrow \langle \pi_\ell^{**}(x, b'), a \rangle = \langle b', b \rangle.$$

It follows that $\pi_\ell^*(b', e_\alpha) \overset{w^*}{\to} b'$ in B^*. Let $b'' \in B^{**}$ and $(b_\beta)_\beta \subseteq B$ such that $b_\beta \overset{w^*}{\to} b''$ in B^{**}. Since $Z^t_{e''}(B^{**}) = B^{**}$, for every $b' \in B^*$, we have the following equality

$$\langle \pi_\ell^{***}(e'', b''), b' \rangle = \lim_\alpha \lim_\beta \langle b', \pi_\ell(e_\alpha, b_\beta) \rangle$$

$$= \lim_\beta \lim_\alpha \langle b', \pi_\ell(e_\alpha, b_\beta) \rangle = \lim_\beta \langle b', b_\beta \rangle$$

$$= \langle b'', b' \rangle.$$

It follows that $\pi_\ell^{***}(e'', b'') = b''$, and so e'' is a left unit for B^{**}.
Conversely, let e'' be a left unit for B^{**} and suppose that $b \in B$. Thren for every $b' \in B^*$, we have

$$\langle b', \pi(e_\alpha, b) \rangle = \langle \pi^{***}(e_\alpha, b), b' \rangle = \langle e_\alpha, \pi^{**}(b, b') \rangle = \langle \pi^{**}(b, b'), e_\alpha \rangle$$

$$= \langle e'', \pi^{**}(b, b') \rangle = \langle \pi^{***}(e'', b), b' \rangle = \langle b', b \rangle.$$

Then we have $\pi_\ell^*(b', e_\alpha) \overset{w}{\to} b'$ in B^*, and so by Cohen factorization theorem we are done.

$\square$

Corollary 4-14. Let B be a Banach left $A-module$ and A has a $LBAI$ $(e_\alpha)_\alpha \subseteq A$ such that $e_\alpha \overset{w^*}{\to} e''$ in A^{**} where e'' is a left unit for A^{**}. Suppose that $Z^t_{e''}(B^{**}) = B^{**}$. Then $\pi_\ell^*(b', e_\alpha) \overset{w}{\to} b'$ in B^* if and only if e'' is a left unit for B^{**}.

Problem. Let B be a Banach $A-bimodule$. Then
i) If B factors, when B^* factors?
ii) If B is separable, dose B necessarily factor on the one side?

5. Involution $*-$algebra and Arens regularity of module actions

Definition 5-1. Let B be a Banach left $A-module$ and let $(a_\alpha)_\alpha \subseteq A$ has *weak** limit in A^{**}. We say that $a_\alpha)_\alpha$ is left regular with respect to B, if for every $(b_\beta)_\beta \subseteq B$,

$$w^* - \lim_\alpha w^* - \lim_\beta a_\alpha b_\beta = w^* - \lim_\beta w^* - \lim_\alpha a_\alpha b_\beta,$$

where $(b_\beta)_\beta$ has *weak** limit in B^{**}.
The definition of the right regular is similar. For a Banach $A-bimodule$ B, if $(a_\alpha)_\alpha \subseteq A$ is left and right regular with respect to B, we say that $(a_\alpha)_\alpha$ is regular with respect to B. If $(a_\alpha)_\alpha$ is left or right regular with respect to A, we write $(a_\alpha)_\alpha$ is left or right regular, respectively.

Example 5-2.
i) Let B be a right Banach $A-module$ and $a'' \in A^{**}$. Suppose that $Z^r_{a''}(B^{**}) = B^{**}$ and $w^* - \lim_\alpha = a''$ where $(a_\alpha)_\alpha \subseteq A$. Then $(a_\alpha)_\alpha$ is right regular with respect to B.

ii) Let A be a Banach algebra and $(a_\alpha)_\alpha \subseteq A$ *weak** convergence to some point of $Z_1(A^{**})$. Then it is clear that $(a_\alpha)_\alpha$ is regular.

Theorem 5-3. i) Let B be a Banach left (resp. right) $A-module$ and suppose that $(e_\alpha)_\alpha \subseteq A$ is a $LBAI$ for B. If $(e_\alpha)_\alpha \subseteq A$ is left (resp. right) regular with respect to B, then B^* factors on the left (resp. right).
ii) Let B be a Banach left (resp. right) $A-module$ and suppose that B^{**} has a left (resp. right) unit as A^{**}-module. If B^*A (resp. AB^*) is closed subspace of B^*, then B^* factors on the left (resp. right).

Proof. Let $e'' \in A^{**}$ and $e_\alpha \overset{w^*}{\to} e''$. Assume that $(b_\beta)_\beta \subseteq B$ such that $b_\beta \overset{w^*}{\to} b''$ in B^{**}. Then

$$e''b'' = w^* - \lim_\alpha e_\alpha b'' = w^* - \lim_\alpha w^* - \lim_\beta e_\alpha b_\beta = w^* - \lim_\beta w^* - \lim_\alpha e_\alpha b_\beta$$
$$= w^* - \lim_\beta b_\beta = b''.$$

Thus for every $b' \in B^*$, we have

$$\langle b'', b' \rangle = \langle e''b'', b' \rangle = \lim_\alpha \langle e_\alpha b'', b' \rangle = \lim_\alpha \langle b'', b'e_\alpha \rangle.$$

It follows that $b'e_\alpha \overset{w^*}{\to} b'$, and so by Cohen factorization theorem, we are done.
ii) Assume that $B^*A \neq B^*$. Let $e'' \in A^{**}$ be a left unit element for B^{**} and suppose that there is a net $(e_\alpha)_\alpha \subseteq A$ such that $e_\alpha \overset{w^*}{\to} e''$. By Hahn Banach theorem take $0 \neq b'' \in B^{**}$ such that $\langle b'', B^*A \rangle = 0$. Then for every $b' \in B^*$, we have

$$\langle b'', b' \rangle = \langle e''b'', b' \rangle = \lim_\alpha \langle e_\alpha b'', b' \rangle = \lim_\alpha \langle b'', b'e_\alpha \rangle.$$

That is contradiction. Thus $B^*A = B^*$.
Proof of the next part is similar. □

Definition 5-4. Let B be a Banach $A-bimodule$ and suppose that A is a Banach $*-involution$ algebra. We say that B is a Banach $*-involution$ algebra as $A-module$, if the mapping $b \to b^*$ from B into B satisfies in the following conditions

$$(ab)^* = b^*a^*,\ (ba)^* = a^*b^*,\ (\lambda b)^* = \bar{\lambda} b^*,\ (b^*)^* = b\ ,\ \| b^* \|,$$

for all $a \in A$, $b \in B$ and $\lambda \in \mathbb{C}$.

Theorem 5-5. Let B be a Banach $A-bimodule$ and suppose that $(e_\alpha)_\alpha \subseteq A$ is a $LBAI$ for B. Let $(e_\alpha)_\alpha \subseteq A$ be a left regular with respect to B. If B^{**} is a Banach $*-involution$ algebra as A^{**}-module, then B^{**} is unital as A^{**}-module and B^* factors on the both side.

Proof. By using Theorem 5-3, B^* factors on the left and without loss generally, there is a left unit for B^{**} as $e'' \in A^{**}$ such that $e_\alpha \overset{w^*}{\to} e''$. Let $b'' \in B^{**}$. Then

$$b''(e'')^* = (e''(b'')^*)^* = ((b'')^*)^* = b''.$$

Since $e'' = (e'')^*$, B^{**} is unital. Thus it is similar to Theorem 5-3, B^* factors on the right. $\square$

Corollary 5-6. Let B be a Banach $A-bimodule$ and suppose that $(e_\alpha)_\alpha \subseteq A$ is a BAI for B. Then if B^{**} is a Banach $*-involution$ algebra as A^{**}-module, then B^{**} is unital as A^{**}-module and B^* factors on the both side.

Example 5-7. Let G be a locally compact group. Then, on $L^1(G)$ there is a natural involution $*$ defined by $f^*(x) = \Delta(x^{-1})\overline{f(x^{-1})}$, where Δ is modular function and $x \in G$. By preceding theorem if $L^1(G)^{**}$ is a Banach $*-involution$ algebra, then $L^1(G)^{**}$ is unital with respect to first Arens product, and so $LUC(G) = L^\infty(G)$. It follows that G is discrete.

Problem . Let G be a locally compact group. We know that $M(G)$ by $\mu^*(f) = \int \overline{f(x^{-1})}d\mu$ for all $f \in C_0(G)$ is $*$-involution algebra. When there is extension of it into $M(G)^{**}$ where $M(G)^{**}$ equipped first Arens product.

References

1. R. E. Arens, *The adjoint of a bilinear operation*, Proc. Amer. Math. Soc. **2** (1951), 839-848.
2. N. Arikan, *A simple condition ensuring the Arens regularity of bilinear mappings*, Proc. Amer. Math. Soc. **84** (4) (1982), 525-532.
3. J. Baker, A.T. Lau, J.S. Pym *Module homomorphism and topological centers associated with weakly sequentially compact Banach algebras*, Journal of Functional Analysis. **158** (1998), 186-208.
4. F. F. Bonsall, J. Duncan, *Complete normed algebras*, Springer-Verlag, Berlin 1973.
5. H. G. Dales, A. Rodrigues-Palacios, M.V. Velasco, *The second transpose of a derivation*, J. London. Math. Soc. **2** 64 (2001) 707-721.
6. H. G. Dales, *Banach algebra and automatic continuity*, Oxford 2000.
7. N. Dunford, J. T. Schwartz, *Linear operators.I*, Wiley, New york 1958.
8. M. Eshaghi Gordji, M. Filali, *Arens regularity of module actions*, Studia Math. **181** 3 (2007), 237-254.
9. M. Eshaghi Gordji, M. Filali, *Weak amenability of the second dual of a Banach algebra*, Studia Math. **182** 3 (2007), 205-213.
10. H. Farhadi, F.Ghahramani, *Involutions on the second duals of group algebras and a multiplier problem*, Proceedings of the Edinburgh Mathematical Society **50** (2007), 153-161.
11. E. Hewitt, K. A. Ross, *Abstract harmonic analysis*, Springer, Berlin, Vol I 1963.
12. E. Hewitt, K. A. Ross, *Abstract harmonic analysis*, Springer, Berlin, Vol II 1970.
13. A. T. Lau, V. Losert, *On the second Conjugate Algebra of locally compact groups*, J. London Math. Soc. **37** (2)(1988), 464-480.
14. A. T. Lau, A. *Ülger*, *Topological center of certain dual algebras*, Trans. Amer. Math. Soc. **348** (1996), 1191-1212.
15. S. Mohamadzadih, H. R. E. Vishki, *Arens regularity of module actions and the second adjoint of a derivation*, Bulletin of the Australian Mathematical Society **77** (2008), 465-476.
16. M. Neufang, *Solution to a conjecture by Hofmeier-Wittstock*, Journal of Functional Analysis. **217** (2004), 171-180.

17. M. Neufang, *On a conjecture by Ghahramani-Lau and related problem concerning topological center*, Journal of Functional Analysis. **224** (2005), 217-229.
18. J. S. Pym, *The convolution of functionals on spaces of bounded functions*, Proc. London Math Soc. **15** (1965), 84-104.
19. A. *Ülger*, *Arens regularity sometimes implies the RNP*, Pacific Journal of Math. **143** (1990), 377-399.
20. A. *Ülger*, *Some stability properties of Arens regular bilinear operators*, Proc. Amer. Math. Soc. (1991) **34**, 443-454.
21. A. *Ülger*, *Arens regularity of weakly sequentialy compact Banach algebras*, Proc. Amer. Math. Soc. **127** (11) (1999), 3221-3227.
22. A. *Ülger*, *Arens regularity of the algebra* $A\hat{\otimes}B$, Trans. Amer. Math. Soc. **305** (2) (1988) 623-639.
23. P. K. Wong, *The second conjugate algebras of Banach algebras*, J. Math. Sci. **17** (1) (1994), 15-18.
24. N. J. Young *Theirregularity of multiplication in group algebra* Quart. J. Math. Soc., **24** (2) (1973), 59-62.
25. Y. Zhing, *Weak amenability of module extentions of Banach algebras* Trans. Amer. Math. Soc. **354** (10) (2002), 4131-4151.

Department of Mathematics, University of Mohghegh Ardabili, Ardabil, Iran
Email address: haghnejad@aut.ac.ir